

NÖROGENETİK ÇALIŞMA GRUBU 2012 YILI YAYINLARI

Lahut S, Ömür Ö, Uyan Ö, Ağım ZS, Özoğuz A, **Parman Y, Deymeer F, Oflazer P, Koç F, Özçelik H, Auburger G, Başak AN**. ATXN2 and its neighbouring gene SH2B3 are associated with increased ALS risk in the Turkish population. PLoS One. 2012;7(8):e42956.

Gungor-Tuncer O, **Baykan B**, Altindag E, **Bebek N**, Gurses C, Gokyigit A. Prevalence and characteristics of visual aura in idiopathic generalized epilepsy. Epilepsy Behav. 2012;25(4):573-6.

Steffens M, Leu C, **Caglayan H, Yapici Z**, Yalcin DA, **Baykan B, Bebek N**. Genome-wide association analysis of genetic generalized epilepsies implicates susceptibility loci at 1q43, 2p16.1, 2q22.3 and 17q21.32. Hum Mol Genet 2012;15;21(24):5359-72.

Yavuz EN, Ozdemir O, Catal S, Bebek N, Ozbek U, Baykan B. Bromodomain-containing protein 2 gene in photosensitive epilepsy. Seizure. 2012 Oct;21(8):646-8.

Schubert J, Paravidino R, Becker F, Berger A, **Bebek N**, Bianchi A, et al. Hum Mutat. PRRT2 mutations are the major cause of benign familial infantile seizures. Hum Mutat. 2012;33(10):1439-43.

EPICURE Consortium, Leu C, de Kovel CG, Zara F, Covanis A, Suls A, Weckhuysen S, Kaneva R, **Caglayan H**, Turkdogan D, **Baykan B, Bebek N, Ozbek U**, et al. Genome-wide linkage meta-analysis identifies susceptibility loci at 2q34 and 13q31.3 for genetic generalized epilepsies. Epilepsia 2012;53(2):308-18.

Zimoń M, Baets J, Almeida-Souza L, De Vriendt E, Nikodinovic J, **Parman Y, Battaloğlu E**, et al. Loss-of-function mutations in HINT1 cause axonal neuropathy with neuromyotonia. Nat Genet. 2012 Oct;44(10):1080-3.

Benbir G, Hanagasi F, Ertan S, **Hanagasi HA**, Ertan T. Pure gait disturbance displayed by malingers: Case report of two patients. Clin Neurol Neurosurg. 2012 Nov 2. doi:pii: S0303-8467(12)00523-9. 10.1016/j.clineuro.2012.10.005.

Gezen-Ak D, Dursun E, Bilgiç B, **Hanagasi H**, Ertan T, Gürvit H, Emre M, Eker E, Ulutin T, Uysal O, Yilmazer S. Vitamin d receptor gene haplotype is associated with late-onset Alzheimer's disease. Tohoku J Exp Med. 2012;228(3):189-96.

Guerreiro RJ, Lohmann E, Brás JM, Gibbs JR, Rohrer JD, Gurunlian N, Dursun B, Bilgiç B, **Hanagasi H**, Gurvit H, Emre M, Singleton A, Hardy J. Using Exome Sequencing to Reveal Mutations in TREM2 Presenting as a Frontotemporal Dementia-like Syndrome Without Bone Involvement. Arch Neurol. 2012;t 8:1-7.

Bilgiç B, **Hanagasi HA**, Emre M. Dementia and behavioral neurology: recent advances. J Neurol. 2012;259(5):1006-10.

Lohmann E, Guerreiro RJ, Erginel-Unaltuna N, Gurunlian N, Bilgic B, Gurvit H, **Hanagasi HA**, Luu N, Emre M, Singleton A. Identification of PSEN1 and PSEN2 gene mutations and variants in Turkish dementia patients. *Neurobiol Aging*. 2012;33(8):1850.e17-27.

Bilgic B, Bayram A, Arslan AB, **Hanagasi H**, Dursun B, Gurvit H, Emre M, Lohmann E. Differentiating symptomatic Parkin mutations carriers from patients with idiopathic Parkinson's disease: contribution of automated segmentation neuroimaging method. *Parkinsonism Relat Disord*. 2012;18(5):562-6.

Guerreiro RJ, Lohmann E, Kinsella E, Brás JM, Luu N, Gurunlian N, Dursun B, Bilgic B, Santana I, **Hanagasi H**, Gurvit H, Gibbs JR, Oliveira C, Emre M, Singleton A. Exome sequencing reveals an unexpected genetic cause of disease: NOTCH3 mutation in a Turkish family with Alzheimer's disease. A homozygous frameshift mutation of sepiapterin reductase gene causing parkinsonism with onset in childhood. *Neurobiol Aging*. 2012;33(5):1008.e17-23.

Lohmann E, Köroğlu Ç, **Hanagasi HA**, Dursun B, Taşan E, Tolun A. *Parkinsonism Relat Disord*. 2012;18(2):191-3.

Lohmann E, **Hanağası HA**, Gürvit H, Kayacı G, Ruberg M, Yazıcı J, Emre M. The impact of familial structure on Parkinson's disease in Istanbul Medical School, Turkey. *Int J Neurosci*. 2012;122(2):102-5.

Böhm J, Biancalana V, Dechene ET, Bitoun M, Pierson CR, Schaefer E, **Karasoy H**, et al. Mutation spectrum in the large GTPase dynamin 2, and genotype-phenotype correlation in autosomal dominant centronuclear myopathy. *Hum Mutat*. 2012;33(6):949-59

Salar S, Yeni N, Gündüz A, Güler A, Gökçay A, Veliöğlu S, Gündoğdu A, **Hande Çağlayan S**. Four novel and two recurrent NHLRC1 (EPM2B) and EPM2A gene mutations leading to Lafora disease in six Turkish families. *Epilepsy Res*. 2012;98(2-3):273-6.

Tolun A, Scarbrough PM, Zhang H, McKillop JA, Wang F, Kishnani PS, Millington DS, Young SP, Il'yasova D. Systemic oxidative stress, as measured by urinary allantoin and F(2)-isoprostanes, is not increased in Down syndrome. *Ann Epidemiol*. 2012;22(12):892-4.

Güven A, Gündüz A, Bozoglu TM, Yalcinkaya C, **Tolun A**. Novel NDE1 homozygous mutation resulting in microhydranencephaly and not microlyssencephaly. *Neurogenetics*. 2012 Aug;13(3):189-94.

Clark PJ, Thompson AJ, Vock DM, Kratz LE, **Tolun A**, Muir AJ, McHutchison JG, Subramanian M, Millington DM, Kelley RI, Patel K. Hepatitis C virus selectively perturbs the distal cholesterol synthesis pathway in a genotype-specific manner. *Hepatology*. 2012;56(1):49-56.

Tolun A, Graham C, Shi Q, Sista RS, Wang T, Eckhardt AE, Pamula VK, Millington DS, Bali DS. A novel fluorometric enzyme analysis method for Hunter syndrome using dried blood spots. *Mol Genet Metab*. 2012;105(3):519-21.

Lohmann E, Köroğlu Ç, Hanagasi HA, Dursun B, Taşan E, **Tolun A**. A homozygous frameshift mutation of sepiapterin reductase gene causing parkinsonism with onset in childhood. *Parkinsonism Relat Disord*. 2012;18(2):191-3.

Thangarajah T, Lakdawala A, **Battaloglu E**, Malik A, Tillu A. Lengthening z-osteotomy of the fibula to correct persistent talar shift following open reduction internal fixation of ankle fractures. *Foot Ankle Spec*. 2012;5(2):107-10.

Bilir B, Yapici Z, Yalcinkaya C, Baris I, Carvalho C, Bartnik M, Ozes B, Eraksoy M, Lupski J, **Battaloglu E**. High frequency of GJA12/GJC2 mutations in Turkish patients with Pelizaeus-Merzbacher disease. *Clin Genet*. 2012; 27. doi: 10.1111/j.1399-0004.

Berber E, Pehlevan F, Akin M, Yalcin Capan O, Kavakli K, **Caglayan SH**. A Common VWF Exon 28 Haplotype in the Turkish Population. *Clin Appl Thromb Hemost*. 2012; 2: [Epub ahead of print] PMID:22473027.

Gökdemir S, **Cağlayan H**, Kızıltan M, Karaağaç N, Leblebici C, Yeni SN. Presentation of an unusual patient with Lafora disease. *Epileptic Disord*. 2012;14(1):94-8.

Sahin K, Tuzcu M, **Basak N**, Caglayan B, Kilic U, Sahin F, Kucuk O. Sensitization of Cervical Cancer Cells to Cisplatin by Genistein: The Role of NFκB and Akt/mTOR Signaling Pathways. *J Oncol*. 2012;2012:461562. doi: 10.1155/2012/461562.

Roy A, **Basak NP**, Banerjee S. Notch1 intracellular domain increases cytoplasmic EZH2 levels during early megakaryopoiesis. *Cell Death Dis*. 2012;23;3:e380.

Gunes S, Asci R, Okten G, Atac F, Onat OE, Ogur G, Aydin O, **Ozcelik T**, Bagci H. Two Males with SRY-Positive 46,XX Testicular Disorder of Sex Development. *Syst Biol Reprod Med*. 2012 Oct 30. [Epub ahead of print].

Emre Onat O, Gulsuner S, Bilguvar K, **Nazli Basak**, Topaloglu H, Tan M, Tan U, Gunel M, Ozcelik T. *Eur J Hum Genet*. 2012 Aug 15. doi: 10.1038/ejhg.2012.170

Sarac O, Gulsuner S, Yildiz-Tasci Y, **Ozcelik T**, Kansu T. Neuro-ophthalmologic findings in humans with quadrupedal locomotion. *Ophthalmic Genet*. 2012;33(4):249-52.

Emre Onat O, Gulsuner S, Bilguvar K, **Nazli Basak A**, Topaloglu H, Tan M, Tan U, Gunel M, **Ozcelik T**. Missense mutation in the ATPase, aminophospholipid transporter protein ATP8A2 is associated with cerebellar atrophy and quadrupedal locomotion. *Eur J Hum Genet*. 2012 Aug 15. doi: 10.1038/ejhg.2012.170.